

*Proceeding of the 5th Congress of Asian Association of Veterinary Anatomists
Bali- INDONESIA, February 11-13th, 2015*

AH-19

The Heavy Metal Deposit Visualization on Organ Tissues after Slaughtering on Beef Production

Adi Winarto^{1,*}, I Ketut Mudite Adnyane², and Yohana Paula Prihatmi Pamungkasari³

^{1,2} Laboratory of Histology, Department of Anatomy, Physiology and Pharmacology,
Faculty of Veterinary Medicine, Bogor Agricultural University

³ Citra Pet & Vet Ruko City. Walk 6 No.12. Citragrand, Cibubur-Bekasi

*Corresponding author: katanablack704@gmail.com

Keywords: lead, heavy metal, cow's tissue, AAS, Rhodizonate

INTRODUCTION

Currently, heavy metal especially lead (Pb) is highly potential in polluting environment that may harmful to animal productions and human health because of its accumulation (Harteman 2011). The lead accumulation effect to the risk organs gating wors by increasing deposit number in the tissues target. Since beef is one of animal protein resources of human meal, it is important to consider the lead accumulation effect to the human health (Suyanto *et al.* 2010). Over Pb accumulation in vital organs has a toxic effect to the individual (D'Mello 2003). This study was aimed to visualize lead deposit at the muscle, liver and kidney tissues and to examine the lead levels in evaluated tissues.

MATERIALS AND METHODS

The tissue samples were collected from slaughter houses surrounds Bogor. Tissue samples were divided into 2 parts, one part was fixed in 4 % paraformaldehyde for histological method, which lead was visualized by *Rhodizonate* (Kiernan 1990), and the other part was prepared for lead value level analysis using Atomic Absorbent Spectrophotometer.

RESULTS AND DISCUSSION

The result showed that the accumulation lead in muscle was scatter in outer connective tissue rather than in muscle cell. In the liver, lead accumulation was found in hepatocyte at the peripheral area of lobules. While in the kidney, lead was mainly deposited in renal tubule, although slightly also found in glomerular area. Further evaluation using AAS indicated that all evaluated organ samples showed an exceed value from recommended maximum number approved by Indonesian government (SNI 7387:2009). Those lead values levels in the muscle, liver, and kidney tissues, respectively were as followed 1.24 ± 0.33 ppm, 1.59 ± 0.61 ppm, 1.11 ± 0.20 ppm. The result indicated that excluded the outer muscle connective tissue in beef production will reduce a risk in accumulating lead deposit, and consuming liver and kidney was potentially catch lead from meal, which harmful to the human health. More over these data may indicate how important the regular control on lead pollutant to improve beef quality.

CONCLUSION

It can be conclude that both approach giving an easy guidance in understanding the risk of getting lead through the "meat" production. The histochemical data indicated that there may get a hepatocyte function failure risk and get a renal tubular damage when the lead accumulation was there.

REFERENCES

1. D'Mello JPF. 2003. *Food Safety Contaminants and Toxins*. London: CABI Publishing.
2. Harteman E. 2011. Dampak kandungan logam berat terhadap kemunculan polimorfisme ikan badik (*Arius maculatus* Fis & Bian) dan sembilang (*Plotosus canius* Web & Bia) di muara

Sungai Kahayan serta Katingan, Kalimantan Tengah[disertasi]. Bogor (ID): Institut Pertanian Bogor. 268 hlm.

3. Kiernan JA. 1990. *Histological and Histochemical Methods*. 2nd ed. Canada: Pergamen Pr.
4. [SNI] Standar Nasional Indonesia 7387:2009. 2009. Batas maksimum cemaran logam berat dalam pangan. Badan Standarisasi Nasional.